

Math 240, Second Exam REVIEW QUESTIONS

1. Find the inverse of the matrix A below:

$$A = \begin{bmatrix} 1 & 1 & -1 \\ 4 & 2 & -1 \\ -2 & -1 & 1 \end{bmatrix}$$

2. Suppose A , B and X are $n \times n$ matrices, with A , X , and $A - AX$ invertible, and suppose

$$(A - AX)^{-1} = X^{-1}B$$

First, explain why B is invertible, then solve the equation for X . If you need to invert a matrix, explain why it is invertible.

3. Let $A = \begin{bmatrix} 1 & 2 \\ 5 & 12 \end{bmatrix}$. Find A^{-1} using the formula, then solve $A\mathbf{x} = [3, 5]^T$.
4. Show that, if AB is invertible, then so is A (assume A, B are $n \times n$). Hint: If AB is invertible, then there is a matrix W so that $ABW = I$.
5. Let S be the parallelogram whose vertices are $(-1, 1)$, $(0, 4)$, $(1, 2)$ and $(2, 5)$. Use determinants to find the area of S .

6. Let $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$, $B = \begin{bmatrix} a + 2g & b + 2h & c + 2i \\ d + 3g & e + 3h & f + 3i \\ g & h & i \end{bmatrix}$, and $C = \begin{bmatrix} g & h & i \\ 2d & 2e & 2f \\ a & b & c \end{bmatrix}$.

If $\det(A) = 5$, find $\det(B)$, $\det(C)$, $\det(BC)$.

7. Assume that A and B are row equivalent, where:

$$A = \begin{bmatrix} 1 & 2 & -2 & 0 & 7 \\ -2 & -3 & 1 & -1 & -5 \\ -3 & -4 & 0 & -2 & -3 \\ 3 & 6 & -6 & 5 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 & 4 & 0 & -3 \\ 0 & 1 & -3 & 0 & 5 \\ 0 & 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- (a) State which vector space contains each of the four subspaces, and state the dimension of each of the four subspaces:
- (b) Find a basis for $\text{Col}(A)$:
- (c) Find a basis for $\text{Row}(A)$:
- (d) Find a basis for $\text{Null}(A)$:
8. Determine if the following sets are subspaces of V . Justify your answers.

$$\bullet H = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix}, a \geq 0, b \geq 0, c \geq 0 \right\}, \quad V = \mathbb{R}^3$$

- $H = \left\{ \begin{bmatrix} a + 3b \\ a - b \\ 2a + b \\ 4a \end{bmatrix}, a, b \text{ in } \mathbb{R} \right\}, V = \mathbb{R}^4$
- $H = \{f : f'(x) = f(x)\}, V = C^1(-\infty, \infty)$
(C^1 is the space of differentiable functions where the derivative is continuous).
- H is the set of vectors in \mathbb{R}^3 whose first entry is the sum of the second and third entries, $V = \mathbb{R}^3$.

9. Prove that, if $T : V \mapsto W$ is a linear transformation between vector spaces V and W , then the range of T , which we denote as $T(V)$, is a subspace of W .
10. Let H, K be subspaces of vector space V . Define $H + K$ as the set below, and see if $H + K$ is a subspace (check all parts of the definition).

$$H + K = \{\mathbf{w} \mid \mathbf{w} = \mathbf{u} + \mathbf{v}, \text{ for some } \mathbf{u} \in H, \mathbf{v} \in K\}$$

11. Let A be an $n \times n$ matrix. Write statements from the Invertible Matrix Theorem that are each equivalent to the statement “ A is invertible”. Use the following concepts, one in each statement: (a) $\text{Null}(A)$ (b) Basis (c) Rank (d) $\det(A)$
12. Is it possible that all solutions of a homogeneous system of ten linear equations in twelve variables are multiples of one fixed nonzero solution? Discuss.
13. Show that $\{1, 2t, -2 + 4t^2\}$ is a basis for P_2 .
14. Let $T : V \rightarrow W$ be a 1 – 1 and linear transformation on vector space V to vector space W . Show that if $\{T(\mathbf{v}_1), T(\mathbf{v}_2), T(\mathbf{v}_3)\}$ are linearly dependent vectors in W , then $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ are linearly dependent vectors in V .
15. Use Cramer’s Rule to solve the system:

$$\begin{array}{rcl} 2x_1 & +x_2 & = 7 \\ -3x_1 & & +x_3 = -8 \\ & x_2 & +2x_3 = -3 \end{array}$$

16. Let $A = \begin{bmatrix} -6 & 12 \\ -3 & 6 \end{bmatrix}$, and $\mathbf{w} = [2, 1]^T$. Is \mathbf{w} in the column space of A ? Is it in the null space of A ?
17. Prove that the column space is a vector space using a very short proof, then prove it directly by showing the three conditions hold.
18. If A, B are 4×4 matrices with $\det(A) = 2$ and $\det(B) = -3$, what is the determinant of the following (if you can compute it): (a) $\det(AB)$, (b) $\det(A^{-1})$, (c) $\det(5B)$ (d) $\det(3A - 2B)$, (e) $\det(B^T)$

19. True or False, and give a short reason:

- (a) If $\det(A) = 2$ and $\det(B) = 3$, then $\det(A + B) = 5$.
- (b) Let A be $n \times n$. Then $\det(A^T A) \geq 0$.
- (c) If A^3 is the zero matrix, then $\det(A) = 0$.
- (d) \mathbb{R}^2 is a two dimensional subspace of \mathbb{R}^3 .
- (e) Row operations preserve the linear dependence relations among the rows of A .
- (f) The sum of the dimensions of the row space and the null space of A equals the number of rows of A .

20. Let the matrix A and its RREF, R_A , be given as below:

$$A = \begin{bmatrix} 1 & 1 & 7 & 2 & 2 \\ 3 & 0 & 9 & 3 & 4 \\ -3 & 1 & -5 & -2 & 3 \\ 2 & 2 & 14 & 4 & 2 \end{bmatrix} \quad R_A = \begin{bmatrix} 1 & 0 & 3 & 1 & 0 \\ 0 & 1 & 4 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

so that the columns of A are $\mathbf{a}_1, \dots, \mathbf{a}_5$.

Similarly, define Z and its RREF, R_Z , as:

$$Z = \begin{bmatrix} 4 & 5 & 3 & 4 \\ 5 & 6 & 5 & -3 \\ 10 & -3 & 9 & -106 \\ 4 & 10 & 2 & 44 \end{bmatrix} \quad R_Z = \begin{bmatrix} 1 & 0 & 0 & -4 \\ 0 & 1 & 0 & 7 \\ 0 & 0 & 1 & -5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Label the columns of Z as $\mathbf{z}_1, \dots, \mathbf{z}_4$.

- (a) Find the rank of A and a basis for the column space of A (use the notation \mathbf{a}_1 , etc.). Similarly, do the same for Z :
- (b) You'll notice that the rank of A is the rank of Z . Here is a row reduction using some columns of A and Z :

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 2 & 4 & 5 & 3 \\ 3 & 0 & 4 & 5 & 6 & 5 \\ -3 & 1 & 3 & 10 & -3 & 9 \\ 2 & 2 & 2 & 4 & 10 & 2 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & 2 & -1 \\ 0 & 1 & 0 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Are the subspaces spanned by the columns of A and Z equal?

- (c) Let \mathcal{B} and be the set of basis vectors used for the column spaces of A found in (a). Find the change of coordinates matrix $P_{\mathcal{B}}$ that changes the coordinates from \mathcal{B} to the standard basis, then find the coordinates of \mathbf{z}_1 with respect to \mathcal{B} (Hint: The second part does not rely on the first).
- (d) Find the coordinates of \mathbf{z}_4 using the basis vectors in $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3$.

21. Short Answer:

- (a) Define the *kernel* of a transformation T :
- (b) Define the *dimension* of a vector space:
- (c) We said that \mathbb{P}_n is isomorphic to \mathbb{R}^{n+1} . What is the isomorphism?
- (d) If C is 4×5 , what is the largest possible rank of C ?
What is the smallest possible dimension of the null space of C ?
- (e) If A is a 4×7 matrix with rank 3, find the dimensions of the four fundamental subspaces of A .
- (f) Show that the coordinate mapping (from n -dimensional vector space V to \mathbb{R}^n) is onto.

22. Let A be $m \times n$ and let B be $n \times p$. Show that $\text{rank}(AB) \leq \text{rank}(A)$. (Hint: Explain why every vector in the column space of AB is in the column space of A).

23. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

- (a) If T is one-to-one, what is the dimension of the range of T ?
- (b) What is the dimension of the kernel of T if T maps \mathbb{R}^n onto \mathbb{R}^m ? Explain.

24. Find the determinant of the matrix A below:

$$A = \begin{bmatrix} 4 & 8 & 8 & 8 & 5 \\ 0 & 1 & 0 & 0 & 0 \\ 6 & 8 & 8 & 8 & 7 \\ 0 & 8 & 8 & 3 & 0 \\ 0 & 8 & 2 & 0 & 0 \end{bmatrix}$$

For questions about 4.7 (if it is on the exam), see the homework assigned in class.