

Review questions, Exam 3

1. What is the ansatz we use for y in: Chapter 6? Section 5.2?
2. Finish the definitions:
 - The Heaviside function, $u_c(t)$:
 - The Dirac δ -function: $\delta(t - c)$ (Note: the Dirac function should be defined as a certain limit)
 - Define the convolution: $(f * g)(t)$
 - A function is of **exponential order** if:
3. Use the *definition* of the Laplace transform to determine $\mathcal{L}(f)$:
 - (a)
$$f(t) = \begin{cases} 3, & 0 \leq t < 2 \\ 6 - t, & t \geq 2 \end{cases}$$
 - (b)
$$f(t) = \begin{cases} e^{-t}, & 0 \leq t < 5 \\ -1, & t \geq 5 \end{cases}$$
4. Check your answers to Problem 3 by rewriting $f(t)$ using the step (or Heaviside) function, and use the table to compute the corresponding Laplace transform.
5. Show that $f(t) = t^3$ is of exponential order. Repeat with $f(t) = \cos(t)$. (HINT: If needed, you may assume that $\ln(t) < t$ for $t > 0$).
6. Write the following functions in piecewise form (thus removing the Heaviside function):
 - (a) $(t + 2)u_2(t) + \sin(t)u_3(t) - (t + 2)u_4(t)$
 - (b) $\sum_{n=1}^4 u_{n\pi}(t) \sin(t - n\pi)$
7. Determine the Laplace transform, using the table:
 - (a) $t^2 e^{-9t}$
 - (b) $e^{2t} - t^3 - \sin(5t)$
 - (c) $t^2 y'(t)$ (in terms of $Y(s)$)
 - (d) $e^{3t} \sin(4t)$
 - (e) $e^t \delta(t - 3)$
 - (f) $t^2 u_4(t)$
8. Find the inverse Laplace transform, using the table:
 - (a) $\frac{2s - 1}{s^2 - 4s + 6}$
 - (b) $\frac{7}{(s + 3)^3}$
 - (c) $\frac{e^{-2s}(4s + 2)}{(s - 1)(s + 2)}$
 - (d) $\frac{3s - 1}{2s^2 - 8s + 14}$
 - (e) $(e^{-2s} - e^{-3s}) \frac{1}{s^2 + s - 6}$
9. For the following differential equations, solve for $Y(s)$ (the Laplace transform of the solution, $y(t)$). Do not invert the transform.
 - (a) $y'' + 2y' + 2y = t^2 + 4t, y(0) = 0, y'(0) = -1$
 - (b) $y'' + 9y = 10e^{2t}, y(0) = -1, y'(0) = 5$
 - (c) $y'' - 4y' + 4y = t^2 e^t, y(0) = 0, y'(0) = 0$

10. Solve the given initial value problems using Laplace transforms:

- (a) $2y'' + y' + 2y = \delta(t - 5)$, zero initial conditions.
- (b) $y'' + 6y' + 9y = 0$, $y(0) = -3$, $y'(0) = 10$
- (c) $y'' - 2y' - 3y = u_1(t)$, $y(0) = 0$, $y'(0) = -1$
- (d) $y'' + 4y = \delta(t - \frac{\pi}{2})$, $y(0) = 0$, $y'(0) = 1$
- (e) $y'' + y = \sum_{k=1}^{\infty} \delta(t - 2k\pi)$, $y(0) = y'(0) = 0$. Write your answer in piecewise form.

11. For the following, use Laplace transforms to solve, and leave your answer in the form of a convolution:

- (a) $4y'' + 4y' + 17y = g(t)$ $y(0) = 0, y'(0) = 0$
- (b) $y'' + y' + \frac{5}{4}y = 1 - u_{\pi}(t)$, with $y(0) = 1$ and $y'(0) = -1$.

12. Short Answer:

- (a) $\int_0^{\infty} \sin(3t)\delta(t - \frac{\pi}{2}) dt =$ _____
- (b) Use Laplace transforms to solve the first order DE, thus finding which function has the Dirac function as its derivative:

$$y'(t) = \delta(t - c), \quad y(0) = 0$$

- (c) Use Laplace transforms to solve for $F(s)$, if

$$f(t) + 2 \int_0^t \cos(t-x)f(x) dx = e^{-t}$$

(So only solve for the transform of $f(t)$, don't invert it back).

- (d) In order for the Laplace transform of f to exist, f must be _____
- (e) Can we assume that the solution to: $y'' + p(x)y' + q(x)y = u_3(x)$ is a power series?
- (f) Is $x = 0$ an ordinary point for the differential equation: $xy'' + 3x^2y' + y = 4$?

13. Let $f(t) = t$ and $g(t) = u_2(t)$.

- (a) Use the Laplace transform to compute $f * g$.
- (b) Verify your answer by computing $f * g$ using the definition of the convolution.

14. If $a_0 = 1$, determine the coefficients a_n so that

$$\sum_{n=1}^{\infty} na_n x^{n-1} + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

Try to identify the series represented by $\sum_{n=0}^{\infty} a_n x^n$.

15. Write the following as a single sum in the form $\sum_{k=2}^{\infty} c_k(x-1)^k$ (with perhaps a few terms in the front):

$$\sum_{n=1}^{\infty} n(n-1)a_n(x-1)^{n-2} + x(x-2) \sum_{n=1}^{\infty} na_n(x-1)^{n-1}$$

16. Characterize ALL (continuous or not) solutions to

$$y'' + 4y = u_1(t), \quad y(0) = 1, y'(0) = 1$$

(Hint: We could have solved this IVP without Laplace transforms. How?)

17. Use the table to find an expression for $\mathcal{L}(ty')$. Use this to convert the following DE into a linear first order DE in $Y(s)$ (do not solve):

$$y'' + 3ty' - 6y = 1, y(0) = 0, y'(0) = 0$$

18. Find the recurrence relation between the coefficients for the power series solutions to the following:

- (a) $2y'' + xy' + 3y = 0, x_0 = 0.$
 (b) $(1 - x)y'' + xy' - y = 0, x_0 = 0$
 (c) $y'' - xy' - y = 0, x_0 = 1$

19. Exercises with the table:

- (a) Use table entries 5 and 14 to prove the formula for 9.
 (b) Show that you can use table entry 19 to find the Laplace transform of $t^2\delta(t - 3)$ (verify your answer using a property of the δ function).
 (c) Prove (using the definition of \mathcal{L}) table entries 12 and 13.
 (d) Prove (using the definition of \mathcal{L}) a formula (similar to 18) for $\mathcal{L}(y'''(t))$.

20. Find the first 5 terms of the power series solution to $e^x y'' + xy = 0$ if $y(0) = 1$ and $y'(0) = -1$.

21. Find the radius of convergence for all of the following, and find the interval of convergence for (b) and (d):

(a) $\sum_{n=1}^{\infty} \sqrt{n}x^n$

(c) $\sum_{n=1}^{\infty} \frac{(-1)^n n^2 (x + 2)^n}{3^n}$

(b) $\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n+1}} (x + 3)^n$

(d) $\sum_{n=1}^{\infty} \frac{(3x - 2)^n}{n5^n}$