

Sample Questions (Chapter 3, Math 244)

1. True or False?

- (a) The characteristic equation for $y'' + y' + y = 1$ is $r^2 + r + 1 = 1$
- (b) The characteristic equation for $y'' + xy' + e^x y = 0$ is $r^2 + xr + e^x = 0$
- (c) The function $y = 0$ is always a solution to a second order linear homogeneous differential equation.
- (d) In using the Method of Undetermined Coefficients, the ansatz $y_p = (Ax^2 + Bx + C)(D \sin(x) + E \cos(x))$ is equivalent to

$$y_p = (Ax^2 + Bx + C) \sin(x) + (Dx^2 + Ex + F) \cos(x)$$

- (e) The operator $T(y) = y' + t^2 y + 1$ is a linear operator (in y).

2. (a) First, solve the DE: $y'' + 4y' + 3y = 0$. (b) Use Cramer's rule to find the constants, if the initial conditions are $y(0) = 1$, $y'(0) = \alpha$.

3. Construct the operator associated with the differential equation: $y' = y^2 - 4$. Is the operator linear? Show that your answer is true by using the definition of a linear operator.

4. What do we need to check in order to see if two functions, y_1, y_2 form a **fundamental set of solutions** to a given second order linear homogeneous DE?

5. If $W(f, g) = t^2 e^t$ and $f(t) = t$, find $g(t)$.

6. If y_1, y_2 form a fundamental set of solutions to: $t^2 y'' - 2y' + (3+t)y = 0$, and if $W(y_1, y_2)(2) = 3$, find $W(y_1, y_2)(4)$.

7. Given that $y_1 = \frac{1}{t}$ solves the differential equation: $t^2 y'' - 2y = 0$, find a fundamental set of solutions using Abel's Theorem.

8. Give the general solution:

- (a) $y'' - 3y' - 10y = 0$
- (b) $y'' + 4y' + 4y = 0$
- (c) $y'' - 4y' + 5y = 0$

9. Suppose the roots to the characteristic equation are as given below. Write the general solution to the DE, and write down what the second order linear homogeneous DE was.

- (a) $r = -2, 3$
- (b) $r = 1, 1$
- (c) $r = 2 \pm 3i$

10. Rewrite the expression in the form $a + ib$: (i) 2^{i-1} (ii) $e^{(3-2i)t}$ (iii) $e^{i\pi}$

11. Write $a + ib$ in polar form: (i) $-1 - \sqrt{3}i$ (ii) $3i$ (iii) -4 (iv) $\sqrt{3} - i$

12. Write each expression as $R \cos(\omega t - \delta)$

- (a) $3 \cos(2t) + 4 \sin(2t)$
- (b) $-\cos(t) + \sqrt{3} \sin(t)$
- (c) $4 \cos(3t) - 2 \sin(3t)$

13. Practice setting up the Spring-Mass model: If you need it, $g \approx 9.8 = \frac{49}{5}$.

- (a) Suppose a mass of 0.01 kg is suspended from a spring, and the damping factor is $\gamma = 0.05$. If there is no external forcing, then what would the spring constant have to be in order for the system to *critically damped? underdamped?*

- (b) A mass of 0.5 kg stretches a spring an additional 0.05 meters to get to equilibrium. (i) Find the spring constant. (ii) Does a stiff spring have a large spring constant or a small spring constant (explain).
- (c) It takes 6 N of force to stretch a certain spring 3 meters. A mass of $\frac{1}{2}$ kg is attached to a spring. (a) Find the spring constant, and (b) if the damping constant is 2, write the differential equation for the motion of the mass.
- (d) Given the model of motion of the mass on a spring is given by

$$\frac{1}{2}u'' + \gamma u' + 8u = 0$$

Find γ so that the spring is *underdamped*, *critically damped*, *overdamped*.

14. Solve. If there are initial conditions, solve for all constants, otherwise, find the general solution.

- (a) $u'' + u = 3t + 4$, $u(0) = 0$, $u'(0) = 0$.
 (b) $u'' + u = \cos(2t)$, $u(0) = 0$, $u'(0) = 0$
 (c) $u'' + u = \cos(t)$, $u(0) = 0$, $u'(0) = 0$ (And please compare to the previous problem).
 (d) $y'' + 4y' + 4y = e^{-2t}$
 (e) $y'' - 2y' + y = te^t + 4$, $y(0) = 1$, $y'(0) = 1$.

15. Find a second order linear differential equation with constant coefficients whose general solution is given by:

$$y(t) = C_1 + C_2e^{-t} + \frac{1}{2}t^2 - t$$

16. Determine the longest interval for which the IVP is certain to have a unique solution (Do not solve the IVP):

$$t(t-4)y'' + 3ty' + 4y = 2 \quad y(3) = 0 \quad y'(3) = -1$$

17. Let $L(y) = ay'' + by' + cy$ for some value(s) of a, b, c .

If $L(3e^{2t}) = -9e^{2t}$ and $L(t^2 + 3t) = 5t^2 + 3t - 16$, what is the particular solution to:

$$L(y) = -10t^2 - 6t + 32 + e^{2t}$$

18. For each DE below, use the Method of Undetermined Coefficients to give the final form of your guess for the particular solution, $y_p(t)$. Do NOT solve for the coefficients.

- (a) $y'' + 3y' = t^3 + t^2e^{-t} + \sin(3t)$
 (b) $y'' + y = t(1 + \sin(t))$
 (c) $y'' - 5y' + 6y = e^{2t}(3t + 4)\sin(t)$
 (d) $y'' + 2y' + 2y = 3e^{-t} + 2e^{-t}\cos(t) + 4e^{-t}t^2\sin(t)$

19. Each equation below exhibits either **beating**, **resonance**, or **neither**. Label each one with B , R , or N .

- (a) $y'' + 3y = \cos(3t)$
 (b) $y'' + 9y = \cos(3t)$
 (c) $y'' + (1.1)^2y = \sin(t)$
 (d) $y'' + 3y = \cos(9t)$

Be sure to remember that separate homework sheet that took the place of Section 3.8.