

### Selected Solutions, Section 6.3

1. Problem 2: To sketch the graph, try first rewriting what is given as a piecewise defined function. The function:

$$(t-3)u_2(t) - (t-2)u_3(t)$$

depends on the value of  $t$ :

- If  $t < 2$ , the function is zero, since  $u_2$  and  $u_3$  are zero.
- If  $2 \leq t < 3$ , then the function is just  $t-3$ , since  $u_3(t)$  is still zero.
- If  $t \geq 3$ , then the function would be  $t-3 - (t-2) = -3+2 = -1$ , since both  $u_2$  and  $u_3$  would simplify to 1.

Now it's easy to graph.

2. Problem 3: Plot  $(t-\pi)^2 u_\pi(t)$ . This is the right half of  $t^2$  shifted  $\pi$  units to the right.
3. Problem 6: Break it up:

$$(t-1)u_1(t) - 2(t-2)u_2(t) + (t-3)u_3(t) = \begin{cases} 0 & \text{if } t < 1 \\ (t-1) & \text{if } 1 \leq t < 2 \\ (t-1) - 2(t-2) & \text{if } 2 \leq t < 3 \\ (t-1) - 2(t-2) + (t-3) & \text{if } t \geq 3 \end{cases}$$

Rewriting this, we get:

$$= \begin{cases} 0 & \text{if } t < 0 \\ t-1 & \text{if } 1 \leq t < 2 \\ -t+3 & \text{if } 2 \leq t < 3 \\ 0 & \text{if } t \geq 3 \end{cases}$$

4. Problem 8: To use the table, we need that:

$$u_1(t)f(t-1) = u_1(t)(t^2 - 2t + 2)$$

so that  $f(t-1) = t^2 - 2t + 2$ . This means that

$$f(t) = (t+1)^2 - 2(t+1) + 2 = t^2 + 2t + 1 - 2t - 2 + 2 = t^2 + 1$$

Now the table entry says:

$$\frac{f(t)}{u_c(t)f(t-c)} \bigg| \frac{F(s)}{e^{-sc}F(s)}$$

so we see that

$$F(s) = \mathcal{L}(t^2 + 1) = \frac{2}{s^3} + \frac{1}{s}$$

so that our final answer is:  $e^{-s} \left( \frac{2}{s^3} + \frac{1}{s} \right)$

We could verify it directly as well by computing

$$\mathcal{L}(u_1(t)(t^2 - 2t + 2)) = \int_1^\infty e^{-st}(t^2 - 2t + 2) dt$$

5. Problem 11: Find the Laplace transform of

$$f(t) = (t-3)u_2(t) - (t-2)u_3(t)$$

We break it up using the linearity of  $\mathcal{L}$ :

- To use the table, we must have:  $(t-3)u_2(t) = f(t-2)u_2(t)$ . Therefore,

$$f(t) = t+2-3 = t-1 \quad F(s) = \frac{1}{s^2} - \frac{1}{s}$$

and the Laplace transform of  $f(t-2)u_2(t)$  is  $e^{-sc}F(s)$ , so overall, we get:

$$e^{-2s} \left( \frac{1}{s^2} - \frac{1}{s} \right)$$

- Similarly, for the second term,  $(t-2)u_3(t) = f(t-3)u_3(t)$ , so

$$f(t) = t + 3 - 2 = t + 1 \quad F(s) = \frac{1}{s^2} + \frac{1}{s}$$

and overall, the Laplace transform of this part is:

$$e^{-3s} \left( \frac{1}{s^2} + \frac{1}{s} \right)$$

Overall, we subtract the two answers:

$$F(s) = e^{-2s} \left( \frac{1}{s^2} - \frac{1}{s} \right) - e^{-3s} \left( \frac{1}{s^2} + \frac{1}{s} \right)$$

6. Problem 14: Find the inverse Laplace transform of

$$F(s) = e^{-2s} \cdot \frac{1}{s^2 + s - 2}$$

This is of the form  $e^{-sc}F(s)$ , so we need to find the inverse transform of  $1/(s^2 + s - 2)$ . We'll do this by Partial Fraction Decomposition. This  $F$  refers to the table, not the original  $F$ :

$$F(s) = \frac{1}{s^2 + s - 2} = \frac{A}{s+2} + \frac{B}{s-1} = -\frac{1}{3} \frac{1}{s+2} + \frac{1}{3} \frac{1}{s-1}$$

and

$$f(t) = -\frac{1}{3}e^{-2t} + \frac{1}{3}e^t$$

So our overall answer is  $u_2(t)f(t-2)$ , or:

$$u_2(t) \left( -\frac{1}{3}e^{-2(t-2)} + \frac{1}{3}e^{t-2} \right)$$

7. Problem 15: Find the inverse Laplace transform of

$$G(s) = \frac{2e^{-2s}(s-1)}{s^2 - 2s + 2}$$

(I changed the notation of the original function so as not to confuse  $F(s)$  in the table with  $F(s)$  in the original question).

We will rewrite this expression, keeping the table in mind:

$$G(s) = 2e^{-2s} \frac{s-1}{s^2 - 2s + 2} = 2e^{-2s} \frac{s-1}{(s-1)^2 + 1} = 2e^{-sc}F(s)$$

We see that, given this  $F(s)$ , then  $f(t) = e^t \cos(t)$  and our overall inverse Laplace transform is:

$$2u_2(t)f(t-2) = 2u_2(t)e^{t-2} \cos(t-2)$$

8. Problem 18: Rewrite as:

$$G(s) = e^{-s} \frac{1}{s} + e^{-2s} \frac{1}{s} - e^{-3s} \frac{1}{s} - e^{-4s} \frac{1}{s}$$

Each of these is in the form  $e^{-sc}F(s)$ , where  $F(s) = 1/s$ , so  $f(t) = 1$ . Notice that  $f(t-k) = 1$  as well, so that the overall inverse is simply:

$$g(t) = u_1(t) + u_2(t) - u_3(t) - u_4(t)$$

9. Problem 27: Done in class.

10. Problem 29: Use the results of Problem 28- If  $f$  is periodic with period  $T$ , then

$$\mathcal{L}(f(t)) = \frac{\int_0^T e^{-st} f(t) dt}{1 - e^{-sT}}$$

In this case,  $T = 2$ , and

$$\int_0^2 e^{-st} f(t) dt = \int_0^1 e^{-st} dt = -\frac{1}{s} e^{-st} \Big|_0^1 = -\frac{e^{-s}}{s} + \frac{1}{s} = \frac{1}{s} (1 - e^{-s})$$

Put this into the formula:

$$\mathcal{L}(f(t)) = \frac{\frac{1}{s}(1 - e^{-s})}{1 - e^{-2s}}$$

This doesn't seem to be the same answer as in Problem 27; However we might note that:

$$1 - e^{-2s} = 1^2 - (e^{-s})^2 = (1 - e^{-s})(1 + e^{-s})$$